

Tsemo Aristide
 Visitor, University of Toronto
 Department of mathematics
 100 St Georges Street
 tsemoaristide@hotmail.com

THE DIFFERENTIAL GEOMETRY OF COMPOSITION SEQUENCES OF DIFFERENTIABLE MANIFOLDS.

Abstract.

Let $F_0 = B, F_1, \dots, F_n$ be a sequence of differentiable manifolds and $G_l, l \geq 1$ a Lie subgroup of the group of diffeomorphisms of F_l , H_l a central subgroup of G_l . We denote by K_l the quotient of G_l by H_l . We suppose also given a locally trivial principal bundle p_{K_l} over F_{l-1} which typical fiber is K_l . In this paper we study the differential geometry of the problem of the existence of a composition sequence of manifolds $M_n \rightarrow M_{n-1} \rightarrow \dots \rightarrow M_0 = B$, where each map $f_l : M_l \rightarrow M_{l-1}$ is a locally trivial differentiable bundle over M_{l-1} which typical fiber is F_l , and such that the image of the transitions functions is contained in G_l . We associated to this problem a tower of gerbes, and we study its differential geometry: that is we define the notion of connective structure and curvature.

Introduction.

Let $F_0 = B, F_1, \dots, F_n$ be a sequence of differentiable manifolds and $G_l, l \geq 1$ a Lie subgroup of the group of diffeomorphisms of F_l , H_l a central subgroup of G_l . We denote by K_l the quotient of G_l by H_l . We suppose also given a principal bundle p_{K_l} over F_{l-1} which typical fiber is K_l . In this paper we study the differential geometry of the problem of the existence of a composition sequence of manifolds $M_n \rightarrow M_{n-1} \rightarrow \dots \rightarrow M_0 = B$, where each map $f_l : M_l \rightarrow M_{l-1}$ is a locally trivial differentiable bundle over M_{l-1} which typical fiber is F_l , and such that the image of the transitions functions is contained in G_k . We can associate to f_l a bundle over M_{l-1} which typical fiber is G_l , we suppose also that the quotient of the restriction of this bundle to a fiber of f_{l-1} by H_l is p_{K_l} . We associated to this problem a tower of gerbes, and we study its differential geometry.

The construction of those sequences of fibrations fits in the general problem of the construction of structures in mathematics. There are essentially two ways to obtain new structures:

One is the completion procedure, which is used generally to create space in which unsolved problems have a solution. It also shows the link between algebra and analysis.

The other essential manner to construct structures in mathematics is the gluing procedure, which is the bridge from analysis to geometry. It is the one

involved in our problem. To solve the gluing problem, that is to find obstructions to define global objects which are defined locally, one needs cohomology theories interpreted geometrically.

More practically, consider a topological manifold M , it is built by gluing open subsets of a vector space say E , to study M , we can for example generalize tools used to study E . For instance, we can define the notion of continuous functions. Can functions defined locally be defined globally ? This problem is solved using a 1-cohomology theory.

The notion of sheaf of functions leads to the notion of locally trivial bundles. which are particular case of sheaves of functions. One can also try to solve the following problem. Can sheaves of bundles defined locally be defined globally? To solve this problem, one needs the notion of stacks defined by Giraud [4]. The obstructions to solve this problem is given by a 2-cohomology theory.

More generally, one can try to define the notion of sheaf of stacks (3-sheaf) which would lead to the notion of a 3-cohomology theories,..., the notion of n -sheaf, which would lead to the notion of a n -dimension cohomology theory.

To define a notion of n -sheaf, one needs to define first, the notion of n -category, which is not well understood nowadays.

The gluing procedure in the 1-dimensional case is the theory of sheaves. It is used to construct Hilbert schemes (see Gottsche [5]).

In the two dimensional case, it has been established by Giraud , it is the theory of stacks.

I. The first lifting problem.

Let B and F be two differentiable manifolds, G a Lie subgroup of diffeomorphisms of F , H a subgroup of G central in G , we denote by K the quotient G/H . We suppose defined on B a locally trivial K -principal bundle. We want to study the differential geometry of the following problem: Classify the locally trivial bundles over B , which typical fiber is F and such that the family of elements g_{ij} which define each trivialization $(U_i, g_{ij}, i, j \in I)$ is included in G . This means that the $Diff(F)$ bundle associated to the bundle has a G -reduction. We can associate also to a solution of our problem a principal bundle p_G over B which typical fiber is G . We suppose moreover that the quotient of p_G by H is p_K . This problem will be called the first lifting problem.

We have the exact sequence $1 \rightarrow H \rightarrow G \rightarrow K \rightarrow 1$. We will assume that the map $G \rightarrow K$ has local sections.

Our problem is equivalent to the following: study the existence of a bundle over B which typical fiber is G , and which quotient by H is p_K . To solve it, we will define a gerbe over B which classifying cocycle is the obstruction of the existence of such a bundle. This problem will be called the principal problem associated to the first lifting problem.

Let's recall now some facts from gerbe theory.

Definition 1.

Let B be a manifold, a sheaf S of categories on B , is a map $U \rightarrow S(U)$, where U is an open set of B , and $S(U)$ category which satisfies the following properties:

- To each inclusion $U \rightarrow V$, there exists a map $r_{U,V} : S(V) \rightarrow S(U)$ such that $r_{U,V} \circ r_{V,W} = r_{U,W}$.

- Gluing conditions for objects,

Consider a covering family $(U_i)_{i \in I}$ of an open set U of B , and for each i , an object x_i of $S(U_i)$, suppose that there exists a map $g_{ij} : r_{U_i \cap U_j, U_j}(x_j) \rightarrow r_{U_i \cap U_j, U_i}(x_i)$ such that $g_{ij}g_{jk} = g_{ik}$, then there exists an object x of $S(U)$ such that $r_{U_i, U}(x) = x_i$

Gluing conditions for arrows,

Consider two objects P and Q of $S(B)$, then the map $U \rightarrow \text{Hom}(r_{U,B}(P), r_{U,B}(Q))$ is a sheaf.

Moreover, if the following conditions are satisfied the sheaf of categories S is called a gerbe

$G1$

There exists a covering family $(U_i)_{i \in I}$ of B such that for each i the category $S(U_i)$ is not empty

$G2$

Let U be an open set of B , for each objects x and y of U , there exists a covering $(U_i)_{i \in I}$ of U such that $r_{U_i, U}(x)$ and $r_{U_i, U}(y)$ are isomorphic

$G3$

Every arrow of $S(U)$ is invertible, and there exists a sheaf A in groups on B , such that for each object x of $S(U)$, $\text{Hom}(x, x) = A(U)$, and this family of isomorphisms commute with the restriction maps.

The sheaf A is called the band of the gerbe S , in the sequel, we will consider only gerbes with commutative band.

Notation.

for a covering family $(U_i)_{i \in I}$ of B , and an object x_i of $S(U_i)$, we denote by $x_{i_1 \dots i_n}^i$ the element $r_{(U_{i_1} \cap \dots \cap U_{i_n}, U_i)}(x_i)$, and by $U_{i_1 \dots i_n}$ the intersection $U_{i_1} \cap \dots \cap U_{i_n}$.

Let endows B with a gerbe S with band A , one can associates the following two Cech cocycle to B . Consider a covering family $(U_i)_{i \in I}$ such that for each i , $S(U_i)$ is not empty, let x_i and x_j be respectively objects of $S(U_i)$ and $S(U_j)$, we consider a map $g_{ij} : x_{ij}^j \rightarrow x_{ij}^i$, the map $t_{ijk} = g_{ij}g_{jk}g_{ki}$ of $\text{Aut}(x_{ijk}^i)$ is the classifying 2 Cech cocycle.

Theorem 1. see [4]

Two gerbes defined over B with band A are isomorphic if and only if their associated 2-cocycles define the same cohomological class. Conversely for each 2-Cech A cocycle c , one can define a gerbe which associated cocycle is c .

We define now on B the sheaf of categories used to solve our problem. For each open set U of B , we denote by $C(U)$ the category which objects are local trivial bundles over U which typical fiber is F , such that the $\text{Diff}(F)$ bundle associated has a G -reduction. This means that the image of the transitions

functions are elements of G . Moreover we suppose that the quotient of this reduction by H is the restriction p_{KU} , of p_K to U . A map between two objects of $C(U)$, is an isomorphism of bundle which induces a map between their associated principal bundles which pushes forward to the identity of p_{KU} . This means that it is an isomorphism which preserves and acts on each fiber by an element of H .

The map

$$U \longrightarrow C(U)$$

defines on B a gerbe which band is H' , the sheaf of differentiable H functions defined on B .

Let

$$k_{ij} : U_i \cap U_j \longrightarrow K$$

be the the transitions functions of the bundle p_K associated to the covering family $(U_i)_{i \in I}$.

Consider a map

$$g_{ij} : U_i \cap U_j \longrightarrow G$$

which pushes forward to k_{ij} . The map $g_{ij}g_{jk}g_{ki}$ defines the classifying cocycle c_H of the gerbe C .

We can also defined the principal gerbe C_P associated to our problem, for each open set U of B , we will denote by $C_P(U)$, the set of locally trivial bundles defined on U which typical fiber is G , and which pushes forward to the restriction of p_K to U .

Connections on principal bundles.

Let $\mathcal{K}$, $\mathcal{G}$ and $\mathcal{H}$ be respectively the Lie algebra of K , G and H . A connection on p_K is a 1-form $\omega : TB \rightarrow \mathcal{K}$ such that:

1

For each fundamental vector field A_B defined by the element A of $\mathcal{K}$, we have: $\omega(A_B(x)) = A$,

2

For each element g of K , we have $\omega(g^*(X)) = Ad(g^{-1})(X)$.

The curvature of ω is the 2-form $d\omega + \frac{1}{2}[\omega, \omega]$.

The notion connective structure of the principal gerbe.

Let ω be a connection on p_K , and e_U an element of $C_P(U)$, recall that is a principal bundle over U which typical fiber is G . We denote by $Co(e_U, \omega)$ the family of connections defined on e_U which push forward to the restriction of ω to U . Consider the map $p_1 : \mathcal{G} \rightarrow \mathcal{K}$, and the map $p_2 : e_U \rightarrow p_{K,U}$ where $p_{K,U}$ is the restriction of p_K to U . the fact that the connection a defined on e_U pushes forward to the restriction ω_U of ω to U means that $p_2^*(\omega_U) = p_1(a)$.

The map $U \longrightarrow Co(e_U, \omega)$ is called the connective structure associated to ω .

Supposed that $((U_i)_{i \in I}, g_{ij})$ is a trivialization of the bundle p_K . Let e_i be an element of $C_P(U_i)$ endowed with a connection w_i over the restriction of ω to $(p_K)_{U_i}$, the restriction of p_K to U_i .

There exists a map g_{ij} between e_{ij}^j and e_{ij}^i the respective restrictions of e_j and e_i to U_{ij} ,

Consider the form $w_j - g_{ij}^* w_i$, (it pushes forward to a form a_{ij} defined on TU_{ij} since H is central in G), and e_{ijk}^i the restriction of e_i to U_{ijk} . On U_{ijk} , we have the restriction of the forms $w_j - g_{ij}^* w_i$, $w_k - g_{jk}^* w_j$ and $w_k - g_{ik}^* w_i$, define respectively on e_{ijk}^j , e_{ijk}^k , we can define the form

$$g_{jk}^*(w_j - g_{ij}^* w_i) + (w_k - g_{jk}^* w_j) - (w_k - g_{ik}^* w_i) =$$

$$-(g_{ij} g_{jk})^* w_i + g_{ik}^* w_i = g_{ik}^*(w_i - (g_{ij} g_{jk} g_{ki})^* w_i) = g_{ik}^*(w_i - c_{ijk}^{-1} dc_{ijk} w_i)$$

Proposition 1. *The family $d(a_{ij})$ is a 1-cocycle.*

Proof.

On U_{ijk} , we have $a_{jk} - a_{ik} + a_{ij} = c_{ijk}^{-1} dc_{ijk}$, since $d(c_{ijk}^{-1} dc_{ijk}) = 0$, we obtain the result.

Consider now the curvature $K(w_i)$ of the connection w_i , it is the form $w_i + \frac{1}{2}[w_i, w_i]$. For any other form w'_i over ω_i , let $a_i = w_i - w'_i$, we have $K(w_i) = K(w'_i) + da_i$, this implies that $dK(w_i)$ does not depend of the form over ω_i . To set in a conceptual theory, one as to define the cotangent and tangent spaces of the principal gerbe.

Definition 2.

Let U be an open set of B , an element of the tangent space of $C(U)$ will be a family of vectors fields (X_e) where e is an object of $C(U)$, such that if e and f are objects of $C(U)$, then there exists a map $g : e \rightarrow f$ such that $Tg(X_e) = X_f$. An element of the cotangent of will be a family of forms $(\alpha)_e$ such that for each vector field (X_e) , we have $\alpha_f(X_f) = \alpha_e(X_e)$, one can define in an obvious way $\Lambda^* C$.

A n -tangent field of $C(U)$ $nTC(U)$ will be a family of n -vector fields $((X_e^1, \dots, X_e^n)_{e \in \text{Obj}(C(U))})$ where X_e^i is a vector field tangent to the element e of $C(U)$ such for each object e and f of $C(U)$, there is a map $g : e \rightarrow f$ such that $Tg(X_e^i) = X_f^i$.

one deduce that the family of 3-form $dK(w_i)$ define a form Ω on $3TC$ which takes value in $\mathcal{G}$ this form is called the curvature of the associated gerbe.

II. The generalization.

Let $B = F_0, F_1, \dots, F_n$ be a sequence of manifolds. For each F_i , we consider a Lie subgroup G_i of diffeomorphisms of F_i , and a central subgroup H_i of G_i . Let denote by K_i the quotient of G_i by H_i , we also suppose given locally trivial

principal bundles p_{K_i} over F_{i-1} which typical fiber is K_i . We will denote by $\mathcal{H}_i$, $\mathcal{G}_i$ and $\mathcal{K}_i$ the respective Lie algebras of H_i , G_i and K_i .

We want to study the differentiable geometry of the following problem:

classify sequences $M_n \rightarrow M_{n-1} \rightarrow \dots M_1 \rightarrow M_0 = B$, where each map $f_i : M_{i+1} \rightarrow M_i$ is locally trivial differentiable bundle which typical fiber is F_{i+1} , such that the transitions functions are elements of G_{i+1} , so we can define a locally trivial principal bundle $p_{MG_{ii+1}}$ over M_i which typical fiber is G_{i+1} , and consider is quotient $p_{MK_{ii+1}}$ by H_{i+1} . We suppose that the restriction $p_{FK_{ii+1}}$ of $p_{MG_{ii+1}}$ to a fiber of f_{i-1} (naturally diffeomorphic to F_i) is $p_{K_{i+1}}$. We will call this problem the n -lifting problem.

We will associate to our problem, a tower of gerbes, and cocycles which will be the obstructions to solve it.

First we define on B the gerbe C_1 associated to the first lifting problem defined by the data F_1 , H_1 , G_1 and K_1 .

Let U be an open set of B and $e_1(U)$ an element of $C_1(U)$, we can associate to $e_1(U)$, the gerbe $C_2(e_1)$ which solve the first lifting problem which data are F_2 , G_2 and K_2 , over the basis $e_1(U)$. Remark that p_{K_2} induces naturally a K_2 -principal bundle p_{e_1} over $e_1(U)$. The quotient of solutions the principal G_2 -bundles associated to the solutions of this first lifting problem by H_2 must be restriction of p_{e_1} .

Suppose that we have defined the gerbes $C_1, C_2(e_1), \dots, C_i(e_{i-1})$, let e_i be an object of $C_i(e_{i-1})$, we define the gerbe $C_{i+1}(e_i)$ which solve the first lifting problem associated to F_{i+1} , G_{i+1} , K_{i+1} with basis e_i . Remark that $p_{K_{i+1}}$ induces naturally a K_{i+1} -principal bundle over e_i , we denote it by p_{e_i} . We suppose that quotient of the principal G_{i+1} bundles associated to the solutions of this first lifting problem by H_{i+1} are restriction of p_{e_i} .

We have assumed that the group H_i is central in G_i , let (U_{jk}^i, k^i_{jk}) be the trivialization of the bundle p_{K_i} , one defines over F_{i-1} a locally trivial bundle Lie_{ii-1} which typical fiber is H_i and which trivialization maps are:

$$U_{jk} \times \mathcal{G}_i \longrightarrow U_{jk} \times \mathcal{G}_i$$

$$(x, y) \longrightarrow (x, Ad(g^i_{jk})(y))$$

where g^i_{jk} is an element of G_i over k^i_{jk} .

this sheaf Lie_{ii-1} is defined for the first lifting problem associated to F_i , H_i , G_i and K_i which basis space is F_{i-1} .

Suppose that we have defined a sheaf over B , $Lie_{ii-1..1}(F_i, H_i, G_i)$ associated to the i -lifting problem defined by the family of groups $F_1, \dots, F_i, H_1, \dots, H_i, G_1, \dots, G_i$. We can define the i -lifting problem associated to the family $F'_1 = F_1, \dots, F'_{i-1} = F_{i-1}, F'_i = Lie_{ii-1}, H'_1 = H_1, \dots, H'_{i-1} = H_{i-1}, H'_i, G_1, \dots, G_{i-1}, G'_i$, where G'_i and H'_i are the groups of automorphisms of Lie_{ii-1} which elements project respectively onto elements of G_i and H_i . We set $Lie_{i+1..1}(F_{i+1}, G_{i+1}, H_{i+1}) = Lie_{i..1}(F'_i, G'_i, H'_i)$.

The classifying cocycle.

We will now define a $n + 1$ -cocycle associated to the tower of gerbes.

For each manifold F_i , we will consider a trivialization $((U^i_j, k^i_{j_1 j_2}, j \in J)$ of the bundle p_{K_i} . Recall that $B = F_0$.

For each open set $U^0_{j_1}$, consider an object $e^1_{j_1}$ of $C_1(U^0_{j_1})$. There exists a map $g^1_{j_1 j_2}$ over $k^1_{j_1 j_2}$ between the respective restrictions of $e^1_{j_2}$ and $e^1_{j_1}$ to $U^0_{j_1 j_2}$.

The map $c^1_{j_1 j_2 j_3} = g^1_{j_1 j_2} g^1_{j_2 j_3} g^1_{j_3 j_1}$ is an automorphism of $e^{1j_1}_{j_2 j_3}$ the restriction of $e^1_{j_1}$ to $U^0_{j_1 j_2 j_3}$ of identified to $U^0_{j_1 j_2 j_3} \times F_1$. the chain $c_{j_1 j_2 j_3}$ can be viewed as a map $U^0_{j_1 j_2 j_3} \rightarrow H_1$. Since C_1 is a gerbe, the family $c^1_{j_1 j_2 j_3}$ is a $2 - H_1$ cocycle.

The map $c^1_{j_1 j_2 j_3}$ acts on the category of open sets of $U^0_{j_1 j_2 j_3} \times F_1$.

The covering which defines the trivialization of $e^1_{j_1 j_2 j_3}$ can be considered as $U^0_{j_1 j_2 j_3} \times U^1$. The map $c_{j_1 j_2 j_3}$ induces a functor $(c_{j_1 j_2 j_3})^*$ between the categories $C_2(U^0_{j_1 j_2 j_3} \times U^1)$ and $C_2(U^0_{j_1 j_2 j_3} \times c_{j_1 j_2 j_3}(U^1))$. Since we have assumed H_1 central in G_1 , we can consider $c_{j_2 j_3 j_4}$, $c_{j_1 j_3 j_4}$, $c_{j_1 j_2 j_4}$ as morphisms of $U^0_{j_1 j_2 j_3} \times F_1$, this enable to compose the functor $(c_{j_1 j_2 j_3})^* (c_{j_2 j_3 j_4})^*$, $(c_{j_1 j_3 j_4})^*$, $(c_{j_1 j_2 j_4})^*$, and to define the Cech boundary $d((c_{j_1 j_2 j_3})^*)$.

Since $c_{j_1 j_2 j_3}$ is a 2-cocycle, the map $d((c_{j_1 j_2 j_3})^*)$ coincide with the restriction of an element $c_{j_1 j_2 j_3 j_4}$ of the set of differentiable functions $U^0_{j_1 j_2 j_3 j_4} \times U^1 \rightarrow H_2$ wich acts by automorphisms on the elements of the category $C_2(U_{j_1 j_2 j_3 j_4} \times U^1)$.

Proposition 1.

The family $c_{j_1 j_2 j_3 j_4}$ defines an 3-Cech cocycle.

Proof.

We have $d(c_{j_1 j_2 j_3 j_4}) = \sum_{i=1}^{i=5} (-1)^i c_{j_1 \dots \hat{j}_i \dots j_5} = \sum_{i=1}^{i=5} (-1)^i \sum_{k=1}^{k=3} (-1)^k (c_{j_1 \dots \hat{j}_k \dots \hat{j}_i \dots})^* = 0$.

Supposed now define the $i + 1$ cocycle associated to the i -lifting problem. It is a family of maps $c_{j_1 \dots j_{i+2}} : U^0_{j_1 \dots j_{i+2}} \times \dots \times U^{i-1} \times F_i$. The map $c_{j_1 \dots j_{i+2}}$ defines a functor $c_{(j_1 \dots j_{i+2})}^*$ between the categories $C_{i+1}(U_{j_1 \dots j_{i+2}} \times U^1 \times \dots \times U^i)$, and the category $C_{i+1}(U_{j_1 \dots j_{i+2}} \times U^1 \times \dots \times U^i c_{j_1 \dots j_{i+2}}(U^i))$, where U^i is an open set of F_i . Since we have supposed that H_i is central in G_i , we can consider the Cech boundary $d(c_{j_1 \dots j_{i+2}})^*$. Since the map $c_{j_1 \dots j_{i+2}}$ is an i -cocycle, we can consider $d(c_{j_1 \dots j_{i+2}})^*$ as a differentiable map $c_{j_1 \dots j_{i+3}} : U_{j_1 \dots j_{i+3}} \times U^1 \times \dots \times U^i \rightarrow H_{i+1}$.

Proposition 2. The family of maps $c_{j_1 \dots j_{i+3}}$ define an $j + 2$ cocycle.

Proof.

We have

$$d(c_{j_1 \dots j_{i+3}}) = \sum_{k=1}^{k=i+4} (-1)^k c_{j_1 \dots \hat{j}_k \dots j_{i+4}} = \sum_{k=1}^{k=i+4} (-1)^k \sum_{l=1}^{l=i+3} (-1)^l (c_{j_1 \dots \hat{j}_l \dots \hat{j}_k \dots j_{i+4}})^* = 0$$

We have seen how to each n -lifting problem one can associate a $n+1$ -cocycle H_n cocycle, is it true that fact conversely to each $n+1-H_n$, one can define a $n+1$ -lifting problem which $n+1$ -classifying cocycle is the given cocycle? if not which conditions must satisfy and $n+1$ -cocycle to be the classifying cocycle of a $n+1$ -lifting problem? We will first study the first lifting problem. Recall that if we have the following exact sequence $1 \rightarrow S_1 \rightarrow S_2 \rightarrow S_3 \rightarrow 1$ of sheaves defined on B such that S_1 is a commutative sheaf, there exists the following exact sequence in cohomology.

$$\begin{aligned} 0 \rightarrow H^0(B, S_1) \rightarrow H^0(B, S_2) \rightarrow H^0(B, S_3) \\ \rightarrow H^1(B, S_1) \rightarrow H^1(B, S_2) \rightarrow H^1(B, S_3) \rightarrow H^2(B, S_3) \end{aligned}$$

denote by δ_1 the map $H^1(B, S_3) \rightarrow H^2(B, S_3)$, we have

Proposition 3.

A $2-H_1$ cocycle c_{H_1} defined on B is the classifying cocycle of a first lifting problem, if and only if there exists an element c_{K_1} of $H^1(B, K_1)$ such that $c_{H_1} = \delta_1(c_{K_1})$.

Proof.

Its enough to remark that each c_{K_1} cocycle c define a principal bundle over B , such that $\delta_1(c)$ is the classifying cocycle of the first lifting problem associated to this bundle.

Remark.

Saying that $\delta_1(c_{K_1}) = c_{H_1}$ means that c_{H_1} is deduced from c_{K_1} in the same way that δ_1 is defined.

Generally the map δ_1 is not injective. To see this consider a G -principal bundle p_G over B , H a central subgroup of G . Denote by K the quotient of G by H and by c the cocycle which defines the quotient p_K of p_G by H , then $\delta_1(c)$ is trivial.

Now suppose that the sheaves S_1 , S_2 and S_3 are commutative, we can write the following long exact sequence

$$\begin{aligned} 0 \rightarrow H^0(B, S_1) \rightarrow H^0(B, S_2) \rightarrow H^0(B, S_3) \\ \rightarrow H^1(B, S_1) \rightarrow H^1(B, S_2) \rightarrow H^1(B, S_3) \rightarrow H^2(B, S_1) \\ \rightarrow \dots \rightarrow H^n(B, S_1) \rightarrow H^n(B, S_2) \rightarrow H^n(B, S_3) \rightarrow H^{n+1}(B, S_3) \end{aligned}$$

denote by δ_n the map $H^n(B, S_3) \rightarrow H^{n+1}(B, S_3)$, we have

Proposition 4.

The $n+1$ -cocycle c_{H_n} of the n -lifting problem is the image of $c_{H_{n-1}}$ by δ_n

Proof.

Straightforward computations.

Definition 1.

We will say that the $n + 1$ -tower is trivial, if and only if its associated cocycle c_{H_n} vanishes.

The principal tower of gerbes associated.

We will define the notion of connective structure and curvature related to the tower of gerbes associated to our problem. First we define the principal tower associated to our problem.

Recall that we can associate to C_1 a principal gerbe C_{1P} defined as follows:

Let U be an open set of B , we consider the category $C_{1P}(U)$ of G_1 principal bundles over U such that the quotient of each of its elements by H_1 is the restriction of p_{K_1} to U ,

Let $e_1(U)$ be an element of $C_1(U)$, we consider an open set U^1 of $e_1(U)$, and an element e_2 of $C_2(U^1)$, it is an F_2 bundle over U^1 , we denote by $p_{U^1} : e_2 \rightarrow U^1$, the canonical projection, we denote by $p_{1U} : U^1 \rightarrow U$ the restriction of the canonical projection of $e_1(U)$ to U^1 .

We can also defined over e_2 the principal bundle e_{2P} associated to the first lifting problem defined by F_2 , H_2 and G_2 which typical fiber is G_2 , the elements of the category $C_{2P}(U^1)$, are fiber product over $p_{1U}(U^1)$ of e_{2P} and e_{1P} , where e_{1P} is a principal bundle over $p_{1U}(U^1)$ associated to the first lifting problem defined by the data F_1 , H_1 and G_1 . Suppose that $U^1 = V \times V^1$ where V and V^1 are open sets of respectively B and F_1 such that the respective restriction of p_{K_1} and p_{K_2} to V and V^1 are trivial. Then the elements of $C_{2P}(U^1)$ are isomorphic to $(V \times G_1) \times (V_1 \times G_2)$. The set of morphisms of elements of $C_{2P}(U^1)$ is given by the natural action of H_2 .

Suppose defined the category $C_{iP}(U^{i-1})$, where U^{i-1} is an open set of an element $C_{i-1}(U^{i-2})$, let U^i be an open set of an object of $C_i(U^{i-1})$, we can define on U^i the principal bundle e_{i+1P} which solve the first lifting problem associated to F_{i+1} , G_{i+1} and H_{i+1} . It is a principal bundle e_{iP} over U^i which typical fiber is G_{i+1} , denote by p_{iU} the canonical map $p_{iU} : U^i \rightarrow U^{i-1}$, the elements of $C_{i+1P}(U^i)$, are fiber product of e_{iP} and e_{i-1P} over $p_{iU}(U^{i-1})$ where e_{i-1P} is an element of $C_{iP}(p_{iU}(U^i))$. The automorphisms group of elements of $C_{i+1P}(U^i)$ are induced by the action of H_{i+1} .

Locally, an object of $C_{iP}(U^{i-1})$ is of the form $U \times G_1 \times (U^1 \times G_2) \times \dots \times (U^{i-1} \times G_i)$.

Definition 1.

The n -tangent space of the tower of gerbes C_P , will be the disjoint unions of the tangent space of the gerbes C_{iP} .

The differential geometry of the principal tower of gerbes.

Suppose defined on each bundle p_{K_i} a connection w_i . We have already define the notion of connective structure for the first lifting problem. Recall that for each element e_1 of $C_{1P}(U)$, it is the family of connections defined on e_1 which project to the restriction of w_1 to the restriction of p_{K_1} to U .

Suppose defined the notion of connective structure for the i -lifting problem, for each object e_{iP} of $C_{iP}(U^{i-1})$, (we can suppose e_{iP} to be isomorphic to $U \times G_1 \times (U^1 \times G_2) \times \dots \times (U^{i-1} \times G_i)$) it is a family of connection defined on e_{iP} , for an element e_{i+1P} of $C_{i+1P}(U^i)$, we will defined the the set $Co(e_{i+1P})$, as the set of $G_1 \times \dots \times G_{i+1}$ connections which project to elements of $Co(e_i(p_{iU}(U^i)))$, where $e_i(p_{iU}(U^i))$ is the canonical projection of e_{i+1P} over an element of $C_{iP}(p_{iU}(U^i))$, consider the projection of U^i to F_i , and the restriction p_{UK_i} of p_{K_i} to the image of the last projection, we suppose also that elements of $Co(e_{i+1P})$ project to the restriction of w_{i+1} .

If this element is isomorphic to $U \times G_1 \times \dots \times U^{i-1} \times G_i$, then an element of $Co(e_{iP})$ is a connection which project to a connection of $Co(e_{i-1P})$, where $e_{i-1P} = U \times G_1 \times \dots \times U^{i-2} \times G_{i-2}$ and which project to a connection of $U^{i-1} \times G_i$, which project to the restriction of w_i to $U^{i-1} \times K_i$.

The sequence of curvatures associated.

For the gerbe associated to the first lifting problem, we have already defined the curvature, it is a $Lie(1)$ 3-form. Let us recall it's definition.

Consider a trivialization $(U_{i_1}, k_{i_1 i_2})$ of the bundle p_{K_1} , over $U_{i_1} \times G_1$ we choose a connection w_{i_1} which project to the restriction of w_1 to $U_{i_1} \times G_1$, we consider the curvature $K(w_{i_1})$ of this connection, then $dK(w_{i_1})$ defines the requested 3-form.

Consider now an element e_i of $C_{iP}(U^{i-1})$, isomorphic to $U_{j_1 \dots j_{i+3}} \times G_1 \times (U^1 \times G_2) \times \dots \times (U^{i-1} \times G_i)$, and $\hat{w}_i$ an element of $Co(e_i)$, if $\hat{w}'_i$ is another element of $Co(e_i)$, then $\hat{w}_i - \hat{w}'_i$ is an $H_1 \oplus \dots \oplus H_n$ form a , we have $K(w_i) = K(w'_i) + da$, this implies that $d(K(w_i))$ is independent of the chosen element of $Co(e_i)$ over w_i

Definition 2.

The form curvature $dK(\hat{w}_i)$, defines a 3-form Ω_i on $3TC_{iP}(U^{i-1})$.

III. Holonomy and parallel displacement.

We will precise the differential geometry of the n -lifting problem by defining notions of holonomy and covariant derivative.

Suppose given a connection ω on p_{K_1} , for each open set U of B and each couple of objects (e_1, e_2) of objects of $C_1(U)$, one can consider connections w_1 and w_2 defined respectively on e_1 and e_2 which push forward to the restriction of ω to the restriction of p_{K_1} to U , unfortunately each isomorphism between e_1 and e_2 does not transforms w_1 to w_2 . To avoid this problem, we are going to enlarge the distribution which defines the connection w_1 and w_2 , to a transitive uniform distribution studied by P. Molino in his thesis, this distribution will behave naturally in respect to the morphisms between objects, generalizing the work of the previous author, will define first, the notion of holonomy and covariant derivative for the principal gerbe C_1 and generalizing those notions for the principal tower associated to the last n -lifting problem. First let recall some results from the work of P. Molino.

Let $p : E \rightarrow B$, be a K -principal locally trivial bundle over B which typical fiber is K , a transitive distribution $\mathcal{D}$ (TD) on E will be a right invariant distribution defined on E such that $d_x p(\mathcal{D}_x) = T_{p(x)} B$.

We will denote by u be the map defined by $u(X) = A$, where X a vertical vector tangent to x and A the element of $\mathcal{K}$ such that $A^*(x) = X$, where A^* is the fundamental vector field defined by A . We will also denote V_x the subset of vertical vector fields at x . The transitive distribution is said to be freely uniform, if $u(\mathcal{D} \cap V_x)$ does not depend of x , this image is thus a normal subgroup L of K .

In the sequel, we will only consider free transitive uniform distributions.

Definition 1.

An horizontal curve of $\mathcal{D}$ is a curve $c : I \rightarrow E$ such that, $c'(t) = \frac{d}{dt}c(t)$ is an element of $\mathcal{D}_{c(t)}$.

Let, x be an element of E , the holonomy group of $\mathcal{D}$ at x , is the set of elements k of K such that x and xk^{-1} can be joined by an horizontal curve.

Let denote by M the quotient of K by L , $U : K \rightarrow M$ the projection map, and $d : \mathcal{K} \rightarrow \mathcal{M}$ the induced Lie algebra morphism, one can define on E , a 1- M form as follows, each element v of $T_x E$ can be written $v = v_1 + v_2$, where v_1 is vertical, and v_2 is an element of $\mathcal{D}_x$, this decomposition is not unique unless that $\mathcal{D}$ is a connection, but the projection $\omega(v)$ of $u(v_1)$ to $\mathcal{M}$ defined on E a 1-form which kernel is $\mathcal{D}$.

Let V be a vector space endowed with a K -action defined by the representation $S : K \rightarrow Gl(V)$, we will denote by $s : \mathcal{K} \rightarrow gl(V)$ the induced Lie algebras representation, by V' the quotient of V by $s(\mathcal{L})$, and by $s' : V \rightarrow V'$ the projection map where $\mathcal{L}$ is the Lie algebra of L . We consider a p -tensorial form $\alpha : TE \rightarrow V$. We denote $\dot{\alpha}$, the V' p -tensorial form defined by $\dot{\alpha}(v_1, \dots, v_p) = s'(\alpha(v_1, \dots, v_p))$ where $v_1, \dots, v_p$ are elements of $T_x E$, write $v_i = w_1^i + w_2^i$, where w_1^i is an vertical vector and w_2^i is an element of $\mathcal{D}_x$ we will define the covariant derivative, $\nabla_{\mathcal{D}}$ of $\dot{\alpha}$ by $\nabla_{\mathcal{D}}(v_1, \dots, v_p) = s'(d\dot{\alpha}(w_2^1, \dots, w_2^p))$.

We have the formula $\nabla_{\mathcal{D}}(\dot{\alpha}) = d\dot{\alpha} + \omega\dot{\alpha}$. We will denote by $\nabla_{\mathcal{D}}\dot{\omega}$, the curvature of $\mathcal{D}$.

We will now apply this notion of TD to the tower of gerbe. First we consider the case of the gerbe C_1 . Let endow p_{K_1} with a connection w_1 and consider an object e_1 of $C_1(U)$, where U is an open set of B . We can define on e_1 the transitive distribution $\mathcal{D}_{1e_1}$, the pull-back of the distribution which define the connection w_1 by the canonical map $e_1 \rightarrow p_{K_1|U}$, if e'_1 is an other object of $C(U)$, then each map $f : e_1 \rightarrow e'_1$ transforms $\mathcal{D}_{1e_1}$ to $\mathcal{D}_{1e'_1}$. The TD $\mathcal{D}_{e_1}$ is uniform, its uniform group is H_1 .

Definition 2.

For each element z of e_1 , we will denote by $Hol(z, e_1, \omega, w_1)$, the holonomy group of the transitive distribution $\mathcal{D}_{e_1}$ in z . Since we have supposed H_1 to be central in G_1 , this group depends only of the projection of z to p_{K_1} .

The holonomy of the tower.

We will define recursively the notion of holonomy of the tower of gerbe, by defining on each element of $C_{i+1P}(U^i)$ a free transitive distribution. We suppose defined on each bundle p_{K_i} a connection w_i .

Let U^1 be an open set of e_1 , recall and object $e_{U^1}^2$ of $C_{2P}(U^1)$, are fibers product over $p_{1U}(U^1)$, of e_{2P} and e_{1P} , where e_{2P} is a principal bundle associated to an object of $C_2(U^1)$ associated to the first lifting problem defined by H_2 , F_2 and G_2 , and e_{1P} is the principal bundle over $p_{1U}(U^1)$ of the first lifting problem associated to the data H_1 , F_1 and G_1 .

We have two canonical projections $f_1 : e_{U^1}^2 \rightarrow e_{1P}$, and $f_2 : e_{U^1}^2 \rightarrow e_{2P}$.

The connection w_2 on p_{K_2} can be pull back to a connection w'_2 on the quotient of e_{2P} by H_2 . Denote by $X_{e_{2P}}$ the pull back to e_{2P} of the distribution which defines the connection w'_2 to e_{2P} , we will define

$$\mathcal{D}_{2e_{U^1}^2} = f_1^* \mathcal{D}_{1e_{1P}} + f_2^* X_{e_{2P}}.$$

Definition 3.

Let z be an element of $e_{U^1}^2$. We denote by $Hol(e_{U^1}^2, \mathcal{D}_2, z, w_1, w_2)$ the holonomy of the transitive distribution $\mathcal{D}_{2e_{U^1}^2}$ in z .

Suppose defined the distribution $\mathcal{D}_{ie_{iP}}$ and the holonomy $Hol(e_{iP}, \mathcal{D}_i, z, w_1, \dots, w_i)$ where is an element of the object e_{iP} , of $C_{iP}(U^{i-1})$.

Recall that an object e^{i+1} of $C_{i+1P}(U^i)$ is a fiber product over $p_{iU}(U^i)$ of e_{iP} and e_{i+1P} , where e_{iP} is an element of $C_{iP}(U^{i-1})$, and e_{i+1P} is a bundle over U^i which solve the first lifting problem associated to F_{i+1} , H_{i+1} and G_{i+1} , we have the canonical projection $f_1 : e^{i+1} \rightarrow e_{iP}$, and $f_2 : e^{i+1} \rightarrow e_{i+1P}$. The connection w_{i+1} on $p_{K_{i+1}}$ induces on the quotient of e_{i+1P} by H_{i+1} a connection ω'_{i+1} . Denote by X_{i+1} the pull back of the distribution which defines the connection ω'_{i+1} to e_{i+1P} , we can define

$$\mathcal{D}_{i+1e^{i+1}} = f_1^*(\mathcal{D}_{ie_{iP}}) + f_2^*(X_{i+1}).$$

Definition 4.

For each element z of e^{i+1} , we denote by $Hol(e^{i+1}, \mathcal{D}_{i+1}, z, w_1, \dots, w_{i+1})$, the holonomy of the transitive distribution $\mathcal{D}_{i+1e^{i+1}}$ in z .

Parallel Displacement.

We will define the notion of parallel displacement for the tower of gerbes.

Definition 5.

Given a connection w_1 on the principal bundle p_{K_1} , and $z : c : I = [0, 1] \rightarrow p_{K_1}$ a differentiable curve, A field X is parallel along c , if and only if $\nabla_{w_1 \frac{d}{dt} c(t)} X(c(t)) = 0$, where ∇_{w_1} is the covariant derivative associated to the connection w_1 .

The parallel displacement along c , is the map $s_{c(t)} : T_{c(0)p_{K_1}} \rightarrow T_{c(1)p_{K_1}}$ such that $s_{c(t)}(v_0)$ is the vector v_1 of $T_{c(1)p_{K_1}}$ such that $v_0 = X(c(0)), v_1 = X(c(1))$ where X is a parallel vector field defined on c , (we suppose c endowed the pull-back of the connection w_1).

Definition 6.

Consider an object e_k of the category $C_{kP}(U^k)$ of the principal tower, and a connection ω of $Co(e_k)$. For each element z of e_k , for each curve $c : I = [0, 1] \rightarrow e_k$, one can define the parallel displacement $s_{c,\omega}$ along c , the parallel displacement along c is the set of maps $s_{c,\omega}, \omega \in Co(e_k)$.

Bibliography.

- [1] Brylinski, J.L Loops spaces, Characteristic Classes and Geometric Quantization, Progr. Math. 107, Birkhauser, 1993.
- [2] Brylinski, J.L, Mc Laughlin D.A, The geometry of degree four characteristic classes and of line bundles on loop spaces I. Duke Math. Journal. 75 (1994) 603-637.
- [3] Deligne, P. Theorie de Hod ge III, Inst. Hautes Etudes Sci. Publ. Math. 44 (1974), 5 – 77.
- [4] Giraud, J. Cohomologie non abélienne. Grundle Math. Wiss. 179, Springer-Verlag.
- [5] Goettsche, L. Hilbert Schemes. Lecture note in Mathematics
- [6] Meinrenken, E. The basic gerbe of a compact simple Lie group. Preprint
- [7] Molino, P. Champs d'éléments sur un espace fibré principal différentiable. Ann. Inst. Fourier 14 (1964) 163-220.
- [8] Murray, M.K., Stevenson, Bundle gerbes: Stable isomorphisms and local theory. J. London. Math. Soc (2) 62 (2000) 925-937
- [9] Perea Medina, A. Flat left-invariant connections adapted to the automorphism structure of a Lie group, J. Differential Geometry, 16 (1981) 445-474.
- [10] Tsemo, A. Non abelian cohomology an application Preprint.